

This is Google's cache of http://www.excedrin.com/8_product_info/8-1a_eselabel.html.

Google's cache is the snapshot that we took of the page as we crawled the web.

The page may have changed since that time. Click here for the [current page](#) without highlighting.

To link to or bookmark this page, use the following url:

http://www.google.com/search?q=cache:SD0qPo2Nen4C:www.excedrin.com/8_product_info/8-1a_eselabel.html+simethicone++microcrystalline+

Google is not affiliated with the authors of this page nor responsible for its content.

These search terms have been highlighted: **simethicone microcrystalline cellulose**

Excedrin® Extra Strength Product Information

Active Ingredients	(per tablet*)	Purpose
Acetaminophen	250 mg	Pain Reliever
Aspirin	250 mg	Pain Reliever
Caffeine	65 mg	Pain Reliever Aid

*available in tablets, caplets & geltabs



INDICATIONS: For temporary relief of the pain of headache, sinusitis, colds, muscular aches, menstrual discomfort, toothache and minor arthritis pain.

DIRECTIONS: Adults: 2 tablets with water every 6 hours while symptoms persist, not to exceed 8 tablets in 24 hours, or as directed by a doctor. Children under 12 years of age consult a doctor.

WARNINGS: Children and teenagers should not use this medicine for chicken pox or flu symptoms before a doctor is consulted about Reye syndrome, a rare but serious illness reported to be associated with aspirin. Keep this and all drugs out of the reach of children. In case of accidental overdose, seek professional assistance or contact a poison control center immediately. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms. As with any drug, if you are pregnant or nursing a baby, seek the advice of a health professional before using this product. **IT IS ESPECIALLY IMPORTANT NOT TO USE ASPIRIN DURING THE LAST 3 MONTHS OF PREGNANCY UNLESS SPECIFICALLY DIRECTED TO DO SO BY A DOCTOR BECAUSE IT MAY CAUSE PROBLEMS IN THE UNBORN CHILD OR COMPLICATIONS DURING DELIVERY.** Do not take this product for pain for more than 10 days or for fever for more than 3 days unless directed by a doctor. If pain or fever persists or gets worse, if new symptoms occur, or if redness or swelling is present, consult a doctor because these could be signs of a serious condition. Consult a

dentist promptly for toothache. Do not take this product if you are allergic to aspirin, have asthma, have stomach problems (such as heartburn, upset stomach or stomach pain) that persist or recur, or if you have ulcers or bleeding problems, unless directed by a doctor. If ringing in the ears or loss of hearing occurs, consult a doctor before taking any more of this product. If you generally consume 3 or more alcohol-containing drinks per day, you should consult your physician for advice on when and how you should take Excedrin and other pain relievers.

DRUG INTERACTION PRECAUTION:

Do not take this product if you are taking a prescription drug for anticoagulation (thinning of blood), diabetes, gout or arthritis unless directed by a doctor.

ACTIVE INGREDIENTS (tablets & caplets): Each tablet or caplet contains:
Acetaminophen 250 mg, Aspirin 250 mg and Caffeine 65 mg.

OTHER INGREDIENTS (tablets & caplets): Benzoic Acid, Hydroxypropylcellulose, Hydroxypropyl Methylcellulose, **Microcrystalline Cellulose**, Mineral Oil, Polysorbate 20, Povidone, Propylene Glycol, **Simethicone** Emulsion, Sorbitan Monolaurate, Stearic Acid. May also contain; Carnauba Wax, FD&C Blue No. 1, Saccharin Sodium, Titanium Dioxide.

ACTIVE INGREDIENTS (geltabs):

Each geltab contains: Acetaminophen 250 mg, Aspirin 250 mg, and Caffeine 65 mg.

OTHER INGREDIENTS (geltabs): Benzoic Acid, D&C Yellow #10 Lake, Disodium EDTA, FD&C Blue #1 Lake, FD&C Red #40 Lake, Ferric Oxide, Gelatin, Glycerin, Hydroxypropylcellulose, Hydroxypropyl Methylcellulose, Maltitol Solution, **Microcrystalline Cellulose**, Mineral Oil, Pepsin, Polysorbate 20, Povidone, Propylene Glycol, Propyl Gallate, **Simethicone** Emulsion, Sorbitan Monolaurate, Stearic Acid, Titanium Dioxide.

Patent Pending (geltabs).

Store at Controlled Room Temperature, 20-25°C (68-77°F).

Questions or comments?

Call 1-800-468-7746.

DISTR. BY: BRISTOL-MYERS PRODUCTS A BRISTOL-MYERS SQUIBB CO.

© 1995, N.Y., N.Y. 10154 MADE IN U.S.A.